

Having trouble getting phosphorus to goal?

XPHOZAH[®]
(tenapanor) tablets

A different way to lower your phosphorus is here.

As add-on therapy for adult patients with chronic kidney disease on dialysis in whom a phosphate binder does not work well:

- ◆ XPHOZAH can help reduce serum phosphorus in adults
- ◆ XPHOZAH is not a binder, it's a blocker
- ◆ XPHOZAH blocks phosphorus throughout the day when taken just before your first and last meal

Ask your healthcare provider about XPHOZAH.



One pill, twice a day.

Scan or visit [XPHOZAH.com](https://www.xphozah.com) to learn more



- ◆ XPHOZAH is dosed as one pill twice a day. If you miss a dose, skip that dose. Do not take 2 doses at the same time
- ◆ Do not take with stool softeners or laxatives, and do not take with a meal right before a hemodialysis session as you may experience diarrhea after taking XPHOZAH. Instead, take before your next meal after dialysis
- ◆ Don't adjust your dose without speaking to your doctor

INDICATION

XPHOZAH (tenapanor) 30 mg BID is a prescription medicine used to reduce serum phosphorus in adults with chronic kidney disease (CKD) on dialysis as add-on therapy when phosphate binders do not work well, or when phosphate binders cannot be tolerated.

IMPORTANT SAFETY INFORMATION

The most important information about XPHOZAH is:

- XPHOZAH may result in softer and/or more frequent bowel movements.
- Do not give XPHOZAH to children who are less than 6 years of age

Please see additional Important Safety Information on next page and full Prescribing Information at [XPHOZAH.com/PI](https://www.xphozah.com/PI).

Lowering phosphorus

Talk with your healthcare provider about your target phosphorus level, and track your progress below.

My target phosphorus level is: _____ mg/dL.

Date	My phosphorus level

IMPORTANT SAFETY INFORMATION (CONT'D)

Do not use XPHOZAH in:

- Children who are less than 6 years of age
- Patients who have a suspected bowel blockage

Before taking XPHOZAH, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed

Do not use XPHOZAH with stool softeners or laxatives. **Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

XPHOZAH can cause serious side effects, including:

Diarrhea is the most common side effect of XPHOZAH, and it can sometimes be severe. Call your doctor if you develop severe diarrhea.

These are not all the possible side effects of XPHOZAH. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to www.fda.gov/medwatch.

For additional safety information, please see full Prescribing Information at XPHOZAH.com/PI.

